



King's Research Portal

DOI:

[10.2217/bmm-2018-0176](https://doi.org/10.2217/bmm-2018-0176)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Kaier, T. E., Alaour, B., & Marber, M. (2018). Cardiac myosin-binding protein C: how a novel biomarker could transform chest pain triage. *Biomarkers in Medicine*, 12(8), 823-826. <https://doi.org/10.2217/bmm-2018-0176>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Editorial: ‘cMyC - how a novel biomarker could transform chest pain triage’

For publication in ‘Biomarkers in Medicine’

Thomas E Kaier, MD, MBA¹; Bashir Alaour, MD¹; Michael Marber, MD, PhD¹;

¹King’s College London BHF Centre, The Rayne Institute, St Thomas’ Hospital, London, UK

Corresponding author: Dr Thomas E Kaier, The Rayne Institute, 4th Floor Lambeth Wing, St

Thomas’ Hospital, Westminster Bridge Road, London SE1 7EH, UK; email: tk@tomkaier.com

Word count: 1,199 (1,500 max)

Keywords:

cMyC; cardiac myosin-binding protein C; cardiac Troponin; chest pain; triage; emergency

department; biomarkers; acute myocardial infarction; AMI

1 **Editorial: ‘cMyC - how a novel biomarker could transform chest pain triage’**

2 Chest pain is a common symptom – according to recent literature it is responsible for at least 6% of
3 all presentations to emergency departments – however, only about 10% of these patients have a
4 final diagnosis of Acute Myocardial Infarction (AMI).[1–5] Chest pain triage is fraught with
5 difficulties as physicians are increasingly caught at the interplay of sensitivity and specificity:
6 Fewer patients now have the diagnostic electrocardiogram (ECG) changes of ST-elevation or
7 depression that allow triage at presentation [6,7] – in fact, 68% of all patients eventually diagnosed
8 with an acute coronary syndrome (ACS) present with Non-ST elevation myocardial infarction
9 (NSTEMI).[8] Consequently, triage has become reliant on the elevation in the blood of the
10 biomarker cardiac Troponin (cTn). This is enshrined in the Universal Definition of Myocardial
11 Infarction [9] and guidelines published by the European Society of Cardiology (ESC) [10],
12 mandating the detection of a cardiac biomarker rise and/or fall for the diagnosis of AMI.

13 **The challenging reality of chest pain triage**

14 The technological advances in evolving cTn assays to high-sensitivity tests comes at the expense of
15 loss of specificity; as analysers are increasingly able to provide quantifiable cTn levels in almost
16 every individual.[11,12] The very definition of a hs-cTn assay – according to the International
17 Federation of Clinical Chemistry and Laboratory Medicine Task Force on Clinical Applications of
18 cardiac Bio-Markers (IFCC TF-CB) – includes 1) a Coefficient of Variation (CV) $\leq 10\%$ at the 99th
19 centile value and 2) the ability to measure at least 50% of healthy individuals with concentrations
20 above the assay’s Limit of Detection (LoD).[13,14] The clinical reality of this advance in assay-

1 technology is that many more patients test ‘Troponin-positive’, but not necessarily ‘AMI-positive’ –
2 all in an attempt to overcome the limitations which made cTn inherently unsuited for early
3 diagnosis of acute myocardial injury: a slow rise and disappearance from the blood stream after
4 myocardial injury.[15,16] By the ESC’s own admission, the clinical implications of using high-
5 sensitivity (hs) cTn assays include a 2-fold increase of detection of type 2 AMI, ~20% relative
6 increase in detection of type 1 AMI and ‘elevations up to 3-fold the upper reference limit
7 (URL)...may be associated with a broad spectrum of conditions’.

8 **Sensitivity comes at a cost**

9 The emergency physician is caught up in this sensitivity/specificity quagmire: they have to handle
10 complex rule-in/rule-out algorithms to optimise care for the patient with suspected Acute Coronary
11 Syndrome (ACS) at the front-door of the hospital.[10,17] Two aspects make this inherently more
12 challenging: 1) Even with high-sensitivity assays, the ESC advocates a delay of 3 hours after chest
13 pain onset for the first blood draw to take place; 2) Many patients get caught up in an ‘observe’
14 zone of indeterminate risk – too high a cTn level for discharge, but too low to classify as AMI.
15 Without doubt, evermore-sensitive assays will establish a new reality of biomarker-interpretation in
16 acute medical services around the world – the (nearly) always-quantifiable level of a cardiac
17 biomarker ought to be scrutinised in the context of the clinical presentation, and we can no longer
18 rely on an outdated black & white approach. But maybe we can achieve more effective triage using
19 a highly-sensitive and specific biomarker with a more favourable release profile?

1 **There might be another way...**

2 Cardiac myosin-binding protein C (cMyC) is a promising novel biomarker of myocardial injury –
3 originally described as the C-protein by Offer et al. in 1973 [18], its discovery relied on the
4 characterisation of ‘impurities’ detected alongside myosin. cMyC has distinctive release-kinetics
5 that should enable it to act as a better adjudicator of acute versus chronic myocardial injury than
6 Troponin.[19] We have raised monoclonal antibodies targeting the cardiac isoform of myosin-
7 binding protein C and successfully migrated the assay onto a high-sensitivity platform.[12]
8 Subsequently, we demonstrated up to 10-fold greater abundance of cMyC after myocardial injury
9 than two leading hs-cTnT/I assays.[20] In a small study involving 174 patients presenting within 3
10 hours of chest pain onset and suspected AMI, we demonstrated a more dynamic rise of cMyC in the
11 early stages of AMI than hs-cTnI.[21] This faster rise ought to yield a positive result (for rule-in of
12 AMI), or an earlier reliable negative result (for rule-out of AMI). Furthermore, the relative
13 abundance of cMyC should allow careful calibration of rule-out and rule-in thresholds, with more
14 ‘headroom’ to enable precise quantification at the low concentrations needed for rule-out.

15 **Large chest pain study confirms efficacy**

16 This hypothesis was tested in a study of >1,900 patients with symptoms suggestive of AMI – in an
17 internal derivation/validation split, we derived cut-off thresholds for immediate rule-out or rule-in
18 of AMI using cMyC instead of hs-cTnT/I (modelled on the 2015 ESC NSTEMI guideline).[10,22]
19 At similar diagnostic accuracy (based on comparable area under the receiver-operating
20 characteristics curve), cMyC was substantially more effective than either hs-cTn assay in guiding

1 patients to (safe) rule-out or rule-in: the net reclassification improvement demonstrated 14.9-23.5%
2 better triage efficiency, thus reducing the size of the ‘observe’ zone substantially. Based on an
3 institution such as St Thomas’ Hospital, a central London hospital home to a tertiary cardiac unit,
4 about 7,800 patients are subject to hs-cTnT testing in the Emergency Department annually. [23] Our
5 findings would translate into savings of 1,000 bed days per year – simply by achieving a more
6 effective triage with a single blood draw at presentation.

7 **Faster, better...Point-of-Care?**

8 But, there is an even greater goal to aim for: point-of-care testing (POCT) of cardiac biomarkers. To
9 date, there is no POCT device that can achieve the levels of sensitivity required to provide accurate
10 measurement of troponin for rule-out of AMI. This task requires a POCT assay to achieve a limit of
11 detection equivalent to the laboratory assay, as the ESC guidelines advocate rule-out only in
12 patients with undetectable hs-cTn levels. The best cTnT POCT platform (Roche Cobas h323
13 handheld instrument) can detect a laboratory-equivalent value of 50 ng/L – about 3.5-fold greater
14 than the 99th centile, or 10-fold the LoD of the laboratory assay.[24] While not bioequivalent, it is
15 tempting to speculate whether cMyC – with a 10-fold greater abundance, and a rule-out threshold
16 25-fold the LoD of the current laboratory platform – might facilitate easier migration onto a
17 handheld device. This would allow rapid deployment of a novel cardiac biomarker to secondary and
18 tertiary care facilities, and pave the way to a cluster-randomised controlled trial. Such a trial, with
19 ethical consent at institutional level to ensure rapid enrolment of a large number of participants,
20 would allow – for the first time – a head-to-head comparison of the effectiveness of different
21 cardiac biomarkers in acute chest pain triage. The potential advantages are compelling: as most

1 patients presenting with chest pain do not have AMI, the goal must be to rule-out AMI in as many
2 patients as possible at the earliest opportunity – i.e. with the first blood draw. Even better, if this
3 could be facilitated in a pre-hospital setting, where cMyC seems to benefit from dynamic release
4 kinetics.[25] cMyC might allow earlier rule-out of AMI [22], and if the promise of more effective
5 triage holds true, up to a fifth of all patients could benefit from expedited discharge, or care where
6 necessary.

7 In conclusion, cMyC is a cardiac-restricted protein which rapidly enters the systemic circulation
8 after myocardial injury and is relatively more abundant than troponin. The biomarker performs
9 favourably in the diagnosis of AMI and is particularly well-suited to a point-of-care diagnostic
10 platform – which could transform the way we perform chest pain triage.

11

1 **Sources of Funding:**

2 The authors are supported by grants from the Medical Research Council (United Kingdom)
3 (G1000737), Guy's and St Thomas' Charity (R060701, R100404), British Heart Foundation
4 (TG/15/1/31518, FS/15/13/31320), and the United Kingdom Department of Health through the
5 National Institute for Health Research Biomedical Research Centre award to Guy's & St Thomas'
6 National Health Service Foundation Trust.

7 Prof Marber is named as an inventor on a patent held by King's College London for the detection of
8 cardiac myosin-binding protein C as a biomarker of myocardial injury.

1 **References**

- 2 1. Goodacre S. The health care burden of acute chest pain. *Heart*. 91(2), 229–230 (2005).
- 3 2. Murphy NF. Hospital discharge rates for suspected acute coronary syndromes between 1990
4 and 2000: population based analysis. *BMJ*. 328(7453), 1413–1414 (2004).
- 5 3. Blatchford O, Capewell S, Murray S, Blatchford M. Emergency medical admissions in
6 Glasgow: general practices vary despite adjustment for age, sex, and deprivation. *Br. J. Gen.*
7 *Pract.* 49(444), 551–554 (1999).
- 8 4. Harris T, McDonald K. Is the case-mix of patients who self-present to ED similar to general
9 practice and other acute-care facilities? *Emerg. Med. J.* 31(12), 970–974 (2014).
- 10 5. Healthcare Quality Improvement Partnership. MINAP Analyses 2012 [Internet]. Available
11 from: [https://data.gov.uk/dataset/a517d8f2-cccd-4f8a-870e-4433bfc1d732/minap-analyses-](https://data.gov.uk/dataset/a517d8f2-cccd-4f8a-870e-4433bfc1d732/minap-analyses-2012)
12 2012.
- 13 6. Jennings SM, Bennett K, Lonergan M, Shelley E. Trends in hospitalisation for acute
14 myocardial infarction in Ireland, 1997-2008. *Heart*. 98(17), 1285–1289 (2012).
- 15 7. Rogers WJ, Frederick PD, Stoehr E, *et al.* Trends in presenting characteristics and hospital
16 mortality among patients with ST elevation and non-ST elevation myocardial infarction in
17 the National Registry of Myocardial Infarction from 1990 to 2006. *Am. Heart J.* 156(6),
18 1026–1034 (2008).
- 19 8. McManus DD, Gore J, Yarzebski J, Spencer F, Lessard D, Goldberg RJ. Recent Trends in
8

the Incidence, Treatment, and Outcomes of Patients with STEMI and NSTEMI. *Am J Med.* 124(1), 40–47 (2011).

9. Thygesen K, Alpert JS, Jaffe AS, *et al.* Third universal definition of myocardial infarction. *Circulation.* 126(16), 2020–2035 (2012).

10. Roffi M, Patrono C, Collet J-P, *et al.* 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Eur. Heart J.* 37(3), 267–315 (2016).

11. Basuray A, French B, Ky B, *et al.* Heart failure with recovered ejection fraction: clinical description, biomarkers, and outcomes. *Circulation.* 129(23), 2380–2387 (2014).

12. Marjot J, Liebetrau C, Goodson RJ, *et al.* The development and application of a high-sensitivity immunoassay for cardiac myosin-binding protein C. *Transl. Res.* 170, 17–25 (2016).

13. Apple FS, Sandoval Y, Jaffe AS, Ordonez-Llanos J, Bio-Markers ITF on CA of C. Cardiac Troponin Assays: Guide to Understanding Analytical Characteristics and Their Impact on Clinical Care. *Clin. Chem.* 63(1), 73–81 (2017).

14. Apple FS, Jaffe AS, Collinson P, *et al.* IFCC educational materials on selected analytical and clinical applications of high sensitivity cardiac troponin assays. *Clin. Biochem.* 48(4–5), 201–203 (2015).

15. Katus HA, Remppis A, Scheffold T, Diederich KW, Kuebler W. Intracellular

1 compartmentation of cardiac troponin T and its release kinetics in patients with reperfused
2 and nonreperfused myocardial infarction. *Am J Cardiol.* 67(16), 1360–1367 (1991).

3 16. Katus HA, Remppis A, Neumann FJ, *et al.* Diagnostic efficiency of troponin T
4 measurements in acute myocardial infarction. *Circulation.* 83(3), 902–912 (1991).

5 17. Amsterdam EA, Wenger NK, Brindis RG, *et al.* 2014 AHA/ACC Guideline for the
6 Management of Patients with Non-ST-Elevation Acute Coronary Syndromes: a report of the
7 American College of Cardiology/American Heart Association Task Force on Practice
8 Guidelines. *J Am Coll Cardiol.* 64(24), e139-228 (2014).

9 18. Offer G, Moos C, Starr R. A new protein of the thick filaments of vertebrate skeletal
10 myofibrils. Extractions, purification and characterization. *J. Mol. Biol.* 74(4), 653–676
11 (1973).

12 19. Baker JO, Tyther R, Liebetrau C, *et al.* Cardiac myosin-binding protein C: a potential early
13 biomarker of myocardial injury. *Basic Res. Cardiol.* , 1–14 (2015).

14 20. Marjot J, Kaier TE, Martin ED, *et al.* Quantifying the Release of Biomarkers of Myocardial
15 Necrosis from Cardiac Myocytes and Intact Myocardium. *Clin. Chem.* 63(5), 990–996
16 (2017).

17 21. Kaier TE, Anand A, Shah AS V, Mills NL, Marber M. Temporal Relationship between
18 Cardiac Myosin-Binding Protein C and Cardiac Troponin I in Type 1 Myocardial Infarction.
19 *Clin. Chem.* 62(8), 1153–1155 (2016).

- 1 22. Kaier TE, Twerenbold R, Puelacher C, *et al.* Direct Comparison of Cardiac Myosin-Binding
2 Protein C With Cardiac Troponins for the Early Diagnosis of Acute Myocardial Infarction.
3 *Circulation*. 136(16), 1495–1508 (2017).
- 4 23. Marjot J, Kaier TE, Henderson K, Hunter L, Marber MS, Perera D. A single centre
5 prospective cohort study addressing the effect of a rule-in/rule-out troponin algorithm on
6 routine clinical practice. *Eur Heart J Acute Cardiovasc Care*. , 2048872617746850 (2017).
- 7 24. Bertsch T, Chapelle J-P, Dempfle C-E, Giannitsis E, Schwabs M, Zerback R. Multicentre
8 analytical evaluation of a new point-of-care system for the determination of cardiac and
9 thromboembolic markers. *Clin. Lab*. 56(1–2), 37–49 (2010).
- 10 25. Kaier TE, Stengaard C, Marjot J, *et al.* Cardiac Myosin-Binding Protein C As Alternative To
11 Cardiac Troponin T For The Diagnosis Of Acute Myocardial Infarction In The Very Early
12 Phase. *J Am Coll Cardiol*. 69(11), 221 (2017).

13